

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets

(11) Publication number:

0 322 016
A1

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 88202800.4

(51) Int. Cl. 4: **C07D 519/00** , **C07D 471/06** ,
C07D 487/06 , **A61K 31/46**

(22) Date of filing: 07.12.88

(30) Priority: 10.12.87 NL 8702981

(43) Date of publication of application:
28.06.89 Bulletin 89/26

(54) Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE

(71) Applicant: **DUPHAR INTERNATIONAL**
RESEARCH B.V.
C.J. van Houtenlaan 36
NL-1381 CP Weesp(NL)

(72) Inventor: Hamminga, Derk
c/o OCTROOIBUREAU ZOAN B.V. P.O. Box
140
NL-1380 AC Weesp(NL)

Inventor: Haeck, Hans H.
c/o OCTROOIBUREAU ZOAN B.V. P.O. Box
140


NL-1380 AC Weesp(NL)
Inventor: van Wijngaarden, Ineke
c/o OCTROOIBUREAU ZOAN B.V. P.O. Box
140

NL-1380 AC Weesp(NL)
Inventor: Wouters, Wouter
c/o OCTROOIBUREAU ZOAN B.V. P.O. Box
140
NL-1380 AC Weesp(NL)

(74) Representative: Muis, Maarten et al
OCTROOIBUREAU ZOAN B.V. P.O. Box 140
NL-1380 AC Weesp(NL)

(54) 1,7-Annulated indolecarboxylic acid esters and -amides.

(57) The invention relates to new 1,7-annulated esters and amides of indolecarboxylic acids having general formula 2,



(2)

wherein the symbols have the meanings given in the specification.


The compounds and their salts are very strong and selective antagonists of "neuronal" 5-hydroxytryptamine (5-HT) receptors, and can be used for the treatment of symptoms which are caused by over-stimulation of these receptors.

EP 0 322 016 A1

1,7-Annulated indolecarboxylic acid esters and -amides.


The invention relates to new 1,7-annulated indolecarboxylic acid esters and -amides of cyclic and polycyclic alcohols or amines, in which cyclic or polycyclic ring a carbon atom has been replaced by a secondary or tertiary nitrogen atom or the N-oxide thereof.

It is known inter alia from European Patent Application 85810595.0 (publication no. 0.189.002) that compounds of the general formula 1



wherein X_1 and X_2 are hydrogen, halogen, alkyl, alkoxy, hydroxy, amino, alkylamino, dialkylamino, mercapto or alkylthio, X_3 is hydrogen, alkyl, alkenyl, aryl or aralkyl, and A is a group -B'-C'-D' which is bound to ring P or ring Q, wherein B' is the group -CO or SO_2 , C' is the group -O- or -NH- and D' is a heterocyclic, nitrogen-containing group, are 5-HT₃ antagonists which may be used for the treatment of gastrointestinal syndromes induced by serotonin.

It has now been found that the new compounds of formula 2



wherein

R_1 is straight or branched alkyl having 1-4 C-atoms; fluorinated or hydroxylated alkyl having 1-4 C-atoms, or R_1 is straight or branched alkenyl or alkynyl having 2-4 C-atoms, or R_1 is straight or branched alkoxy or alkylthio having 1-6 C-atoms which may be substituted with one or more fluorine atoms or hydroxyl groups, or with one optionally substituted phenyl group, or two alkoxy groups R_1 bound to adjacent carbon atoms may form a ring consisting of 5-7 ring atoms, or R_1 is straight or branched (C_{1-4}) alkoxy-, (C_{1-4}) alkylthio- or alkyl- (C_{1-4}) sulphonyl- (C_{1-2}) alkyl which may comprise one or more fluorine atoms or hydroxyl groups, or R_1 is an alkoxycarbonyl methyl group having 1-6 C-atoms in the alkoxy group, or R_1 is a group $R_4R_5N-CO-CH_2-$, R_4R_5N-CO- or $R_4R_5N-SO_2-$ wherein R_4 and R_5 independently of each other are hydrogen, alkyl having 1-3 C-atoms, or together with the nitrogen atom form a heterocyclic 5- or 6-ring, or R_1 is hydroxy, halogen, cyano straight or branched alkoxycarbonyl having 1-6 C-atoms in the alkoxy group;

n has the value 0-3;

Z together with the carbon atom and the nitrogen atom and the intermediate carbon atom forms a heterocyclic group consisting of 5-8 ring atoms, in which, besides the nitrogen atom already present, a second hetero atom from the group N, O, S, S-O, SO_2 or $C=O$ may be present, which ring may be substituted with 1-3 alkyl groups having 1-4 C-atoms, a phenyl group or a spiroalkyl group (C_{2-5}) , or which ring may be annulated with a saturated or non-saturated carbocyclic or heterocyclic ring consisting of 5- or 6-ring atoms which may be substituted with halogen or alkyl having 1-4 C-atoms;

R_3 is hydrogen, halogen, alkyl, alkoxy or alkylthio having 1-4 C-atoms, and

R_2 is the group -B-C-D, wherein B is the group -CO-, C is oxygen or NR_6 , wherein R_6 is hydrogen, (C_{1-7}) alkyl or benzyl, and D is a group $(CH_2)_m-R_7$, wherein m has the value 0, 1 or 2 and R_7 is a carbocyclic ring system in which one carbon atom has been replaced by a secondary or tertiary nitrogen atom or the N-


oxide thereof;

and the pharmacologically acceptable acid addition salts thereof are very strong and selective antagonists of "neuronal" 5-hydroxytryptamine (5-HT) receptors.


The group D is preferably a group of the formula

5


10



(3)



(4)



(5)

wherein p has the value 1 or 2, s has the value 0, 1 or 2, q is 2, 3 or 4, r is 1, 2 or 3, and wherein R_8 is hydrogen, (C_1-7) alkyl, (C_3-8) cycloalkyl, (C_3-8) cycloalkyl- (C_1-2) -alkyl, (C_3-5) alkenyl, (C_3-5) alkynyl, or wherein R_8 is a group $(CH_2)_t-R_9$, wherein t is 1 or 2 and R_9 is hydroxy, or R_9 is the phenyl group, thienyl group, pyrrolyl group or furyl group, which groups may be substituted with 1 or 2 substituents, and R_{10} is absent or is an oxygen atom.

Suitable acids with which the compounds of formula 1 according to the invention can form pharmaceutically acceptable acid addition salts are hydrochloric acid, sulphuric, phosphoric acid, nitric acid, and organic acids, for example, citric acid, fumaric acid, maleic acid, tartaric acid, acetic acid, benzoic acid, *p*-toluenesulphonic acid, methane sulphonic acid and the like.

Both the racemates and (geometric) isomers and the individual enantiomers of compounds of formula 2 belong to the invention.

The antagonistic activity of the compounds of formula 2 on the responses induced by 5-HT or by 2-methyl-5-HT was determined and measured in the Bezold-Jarisch reflex test in rats. The affinity for "neuronal" 5-HT receptors was determined and measured by means of the displacement of (^3H) GR 38032 F of neuroblastoma cells.

On the basis of the antagonistic activity of this type of 5-HT receptors, the compounds may be used for the treatment of symptoms which are caused by over-stimulation of these receptors a) in the gastrointestinal system (nausea and vomiting as a result of exogenic factors, for example, cancer therapy, or endogenic factors, for examples, stasis of the stomach and migraine) ulcer, dyspepsia, spasms, irritable bowel syndrome, etc., or b) in the central nervous system (hallucinations, delusions, manias, anxiety, pain, improvement of vigilance etc.), or c) in the cardiovascular system, for example, spasms of the vessels, arrhythmia, etc, or d) for the relief or prevention of withdrawal syndromes induced for instance by abuse of drugs, or e) in the respiratory system (including nasal disorders or disorders of the bronchi and lungs).


The compounds according to the invention and their salts can be brought into a form suitable for administration, for examples, pills, tablets, coated tablets, capsules, powder, injection liquids and the like by means of techniques conventionally used for this purpose and while using suitable auxiliary substances, for example, solid or liquid carrier materials.

The dosage in which the compounds according to the invention may be used depend on the severity and the nature of the disease to be treated and on the way of administration. As a rule the dosage will be between 0.05 and 20 mg, preferably between 0.1 and 10 mg of active substance daily.

The compounds according to the invention can be prepared in a manner known for analogous compounds, for example:

a) by reaction of a compound of formula 6

50



(6)

55

wherein R_1 , R_3 , n , Z and B have the meanings given in formula 2 and X is a group which may be replaced by a nucleophile, for example, a halogen atom, a 1-imidazolyl group, and alkoxy group, etc., with a

compound of formula 7


$Y - C - D$ (7)

wherein C and D have the meanings given in formula 2 and Y is a hydrogen atom or an alkali metal atom. The reaction is preferably carried out in a suitable solvent, for example, dichloromethane, pyridine, methanol, dimethylformamide, toluene, etc. In particular the compounds according to the invention can be prepared in a good yield



i) by reaction, in the presence of active sodium alkanolate, of a compound of formula 6, wherein R_1 , R_3 , n, Z and B have the meanings mentioned in formula 2 and X is an alkoxy group, with a compound of formula 7 in which C and D have the meanings mentioned in formula 2 and wherein Y is a hydrogen atom. The reaction is preferably carried out in a suitable solvent, for example, toluene, xylene, etc., at temperatures between 20 and 160°C; or

ii) by reaction of a compound of formula 6, wherein R_1 , R_3 , n, Z and B have the meaning given in formula 2 and X is a halogen atom with a compound of formula 7, wherein Y is hydrogen, C represent a group $N-R_6$, and D and R_6 have the meaning given in formula 2. The reaction is preferably carried out in a suitable solvent, for example methylene chloride, acetonitrile, etc. at temperatures between 0 and 80°C.

The starting compounds of formula 6, to be used in these reactions i) and ii) wherein R_1 , R_3 , n, Z and B have the meanings mentioned in formula 2 and X is a hydroxy group or an alkoxy group, may be prepared in a manner known for analogous compounds and may be converted, likewise in known manner, into a compound of formula 6 wherein X is a group which may be replaced by nucleophiles, for example, a halogen atom, a 1-imidazolyl group, etc. Suitable methods of preparing this type of compounds are described in Heterocyclic Compounds (Houlinan, Part 1); J. Am. Chem. Soc. 60, 2414-16 (1938); J.C.S. Perkin Transactions (1981) 636-641. In particular, the starting compounds of formula 8




wherein R_1 , n and Z have the meanings mentioned in formula 2 and R_{11} is alkoxy, can be obtained by reaction of a compound of formula 9



wherein R_1 , n and Z have the meanings mentioned in formula 2. The reaction is preferably carried out in a suitable solvent, for example, tetrahydrofuran, ether, toluene, etc., at temperatures from 0-100°C. The formed reaction product is then cyclised, preferably in the presence of a Lewis acid, for example, zinc chloride, aluminium chloride, etc. The cyclisation is preferably carried out in a suitable solvent, for example, alcohol, benzene, etc. at temperatures between 20 and 150°C.

The compounds of formulae 9 and 10 are known compounds or can be obtained analogously to known compounds.


Starting compounds of formula 11




wherein R_1 , n , Z and R_3 have the meanings given in formula 2, and R_{11} is alkoxy, hydroxy or halogen in particular can be obtained by
 10 i) reaction of a compound of formula 12



wherein R_1 , n and Z have the above-mentioned meanings,
 20 with a compound of formula 13



The reaction is preferably carried out in a solvent, for example alcohol, acetic acid, at temperatures between 20 and 120 °C. The compounds of formula 11 so-obtained wherein R_{11} is alkoxy can be converted in known manner into compounds wherein R_{11} is hydroxy or halogen. The starting compounds having formula 12 and 13 are known compounds, or can be obtained analogous to known compounds; or
 ii) Vilsmeier-Haack formylation of a compound having formula 14



wherein R_1 , n , R_3 and Z have the meanings given in formula 2, followed by oxidation of the formed aldehyde to the desired carboxylic acid, which may be converted in known manner to an analogous starting compound having formula 11. The starting compounds 14 are known or can be prepared analogous to known compound.
 40

b) By substitution of the amine-hydrogen atom in group D in a compound of formula 2, wherein R_1 , R_3 , n , Z , B, C and D have the meanings mentioned in formula 2 with the proviso that the ring nitrogen atom in group D is secondary nitrogen atom, by a group R_8 , wherein R_8 has the above-mentioned meaning, for example, by reaction of a compound of formula 2 with a suitable aldehyde or ketone under reducing conditions, or with a suitable halogen compound. The reaction is preferably carried out in a suitable solvent, for example, water, acetonitrile, alcohol, dimethylformamide, etc. at temperatures between 0 and 150 °C.
 45

The compounds having formula 2 wherein R_{10} is oxygen can be prepared in particular by oxidation of an analogous compound of formula 2 wherein R_{10} is lacking, with a suitable oxidant, for example perbenzoic acid, peracetic acid, hydrogen peroxide, etc., preferably in a suitable solvent, for example, chloroform, acetic acid or water, at temperatures between 0 and 100 °C.
 50

The invention will now be described in greater detail with reference to the ensuing specific examples.

55 EXAMPLE I

(5,6-dihydro-4H-pyrrolo-[3,2,1-ij]-quinoline-1-yl)carboxylic acid (endo-8-methyl-8-azabicyclo-[3,2,1]-oct-3-yl)

ester hydrochloride

0.7 g (30 mmol) of sodium were "dissolved" in 15 ml of methanol. 50 ml of toluene were added, after which the methanol was distilled off, while stirring, until the boiling-point of toluene had been reached, 3.45 g (15 mmol) of (5,6-dihydroxy-4H-pyrrolo-[3,2,1-ij]-quinolin-1-yl)-carboxylic acid ethyl ester and 4.2 g (30 mmol) of tropine were added to this suspension of sodium methanolate in toluene. This mixture was boiled for 4 hours while stirring. After cooling to room temperature, 100 ml of ether and 100 ml of water were added. The organic layer was separated, dried and evaporated to dryness. The residue (5.0 g) was purified by chromatography over 250 g of silica gel. Elution was carried out with a mixture of methanol and triethylamine (97:3). Yield 4.1 g (84%). The ester was converted, by means of alcoholic hydrochloric acid, into the hydrochloride, a substance having a melting-point (with gas formation) of 322 °C.

The following compounds have been prepared in an analogous manner:


- (1) (8-fluoro-5,6-dihydro-4H-pyrrolo-[3,2,1-ij]-quinolin-1-yl)-carboxylic acid-(endo-8-methyl-8-azabicyclo-[3,2,1]-oct-3-yl)ester hydrochloride;
- (2) (5,6-dihydro-4H-pyrrolo-[3,2,1-ij]-quinolin-2-yl)-carboxylic acid-(endo-8-methyl-8-azabicyclo-[3,2,1]-oct-3-yl)ester hydrochloride; yield 58%, melting-point 295-297 °C (decomposition).

20 EXAMPLE IIN-(endo-8-methyl-8-azabicyclo-[3,2,1]-oct-3-yl)-(5,6-dihydro-4H-pyrrolo-[3,2,1-ij]-quinolin-1-yl)-carboxamide hydrochloride

2.0 g (0.01 mol) of 5,6-dihydro-4H-pyrrolo-[3,2,1-ij]-quinolin-1-yl)-carboxylic acid were suspended in 50 ml of methylene chloride. 2.25 ml (0.03 mol) of thionyl chloride were added and the mixture was boiled for 1 hour, and evaporated in vacuo. The residue was dissolved in 50 ml of methylene chloride. 1.5 g (0.01 mol) of 3-amino-8-methyl-8-azabicyclo-[3,2,1]-octane were added, and the mixture was stirred for 30 minutes. Thereafter the reaction mixture was shaken with 2N sodium hydroxide, the methylene chloride layer was washed with water, dried and evaporated in vacuo. The residue was dissolved in ethyl acetate, and a mixture of 0.8 ml of acetyl chloride and 3 ml of alcohol was added. The solid substance was sucked off and dried. Yield 2.9 g (80%) of the desired hydrochloride having a melting-point of 280 °C (decomposition).

The following compounds have been prepared in an analogous manner:

- (1) N-(endo-9-methyl-9-azabicyclo-[3,3,1]-non-3-yl)-5,6-dihydro-4H-pyrrolo-[3,2,1-ij]-quinolin-1-yl)-carboxamide hydrochloride; yield 43%; melting-point 320-323 °C
- (2) N-(endo-8-methyl-8-azabicyclo-[3,2,1]-oct-3-yl)-N-methyl-(5,6-dihydro-4H-pyrrolo-[3,2,1-ij]-quinolin-1-yl)-carboxamide hydrochloride; yield 53%, melting-point 139-141 °C
- (3) N-(1-azabicyclo-[2,2,2]-oct-3-yl)-(5,6-dihydro-4H-pyrrolo-[3,2,1-ij]-quinolin-1-yl)-carboxamide hydrochloride; yield 31%. The product was obtained as a foam. The structure was confirmed by ¹³C-NMR analysis:



15

1	134.22	S	8	122.06	S	15	46.96	T
2	129.33	D	9	24.63	T	16	24.33	T
3	110.22	S	10	22.69	T	17	25.68	D
4	124.42	S	11	44.39	T	18	19.43	T
5	118.69	D	12	165.42	S	19	46.46	T
6	119.33	D	13	45.87	D			
7	121.38	D	14	53.91	T			

20

25

(4) N-(1-azabicyclo-[2,2,2]-oct-3-yl)-(8-fluoro-5,6-dihydro-4H-pyrrolo-[3,2,1-ij]-quinolin-1-yl) carboxamide; yield 52%, melting-point 220-221 °C;

30 (5) N-(1-azabicyclo-[2,2,2]-oct-3-yl)-(4,5,6,7-tetrahydroazepino-[3,2,1-jk]-indol-1-yl)-carboxamide; yield 43%, melting-point 66-69 °C;

(6) N-(endo-8-methyl-8-azabicyclo-[3,2,1]-oct-3-yl)-(4,5,6,7-tetrahydroazepino-[3,2,1-jk]-indol-1-yl)-carboxamide;

35 (7) N-(1-azabicyclo-[2,2,2]-oct-3-yl)-(6,7-dihydro-indolo-[1,7-a,b]-[1]-benzazepin-1-yl)-carboxamide hydrochloride; yield 48%, melting-point 251-252 °C;

(8) N-(1-azabicyclo-[2,2,2]-oct-3-yl)-(6,7-dihydro-indolo-[1,7-ab]-[1]-benzazepin-2-yl)-carboxamide hydrochloride.

40

EXAMPLE III

45 N-{endo-8-(2-hydroxyethyl)-8-azabicyclo-[3,2,1]-oct-3-yl}-(5,6-dihydro-4H-pyrrolo-[3,2,1-ij]-quinolin-1-yl)-carboxamide hydrochloride

(a) 1.16 g (3.6 mmol) of N-(endo-8-methyl-8-azabicyclo-[3,2,1]-oct-3-yl)-(5,6-dihydro-4H-pyrrolo-[3,2,1-ij]-carboxamide were dissolved in 50 ml of 1,2-dichloromethane and cooled to 0 °C under nitrogen. 0.51 g (3.6 mmol) of 1-chloroethyl-chloroformate were added dropwise in 5 min. and the mixture was boiled for 1 hour and evaporated in vacuo. 50 ml of methanol were added to the residue, and the mixture was boiled for 30 min. The mixture was evaporated in vacuo and chromatographed over silicagel with methylene chloride/methanol/ammonia (84/15/1) as eluent. In this manner 0.7 g (63%) of the secondary amine was obtained.

55 (b) This product (2.3 mmol) was dissolved in 35 ml of acetonitrile. 0.18 ml (2.3 mmol) of 2-iodoethanol and 0.63 ml (4.6 mmol) of triethyl amine were added, and the mixture was boiled for 16 hours and then evaporated in vacuo. The residue was shaken with methylene chloride and 2N sodium hydroxide. The methylene chloride layer was dried and evaporated in vacuo. The residue was chromatographed over silicagel with methylene chloride/methanol/ammonia (84/15/1) as eluent. The desired fraction was evap-

orated in vacuo, and the residue was treated with 1.1 equivalent of alcoholic hydrochloric acid. The solid substance was sucked off and dried. In this manner 0.38 g (43%) of the above-mentioned hydrochloride was obtained, having a melting-point of 262-263 °C.

5

EXAMPLE IV


N-(1-azabicyclo-[2,2,2]-oct-3-yl-1-oxide)-(5,6-dihydro-4H-pyrrolo-[3,2,1-ij]-quinolin-1-yl)-carboxamide

10

0.31 g (1 mmol) of N-(1-azabicyclo-[2,2,2]-oct-3-yl)-5,6-dihydro-4H-pyrrolo-[3,2,1-ij]-quinolin-1-yl)-carboxamide were dissolved in 25 ml of chloroform. 0.172 g (1 mmol) of m-chloroperbenzoic acid were added, and the mixture was stirred for 24 hours at room temperature. The mixture was washed with methylene chloride and 2N sodium hydroxide, and evaporated in vacuo. The residue was chromatographed over
15 silicagel. Yield 19%. ¹³C-NMR analysis:

20

25



30

35

40

1	133.74	8	122.22	15	60.06
2	129.09	9	24.76	16	23.39
3	109.04	10	22.26	17	23.77
4	124.33	11	43.85	18	20.38
5	118.51	12	164.54	19	59.53
6	119.04	13	46.38		
7	121.05	14	65.43		

Preparation of intermediates

45

a) 5,6-dihydro-4H-pyrrolo-[3,2,1-ij]-quinolin-1-yl)-carboxylic acid ethyl ester

50

22.5 ml (approximately 0.15 mol) of 90% ethyl bromopyruvate were added dropwise and while stirring in approximately 0.5 hours at a temperature from 20-35 °C to 37.5 ml (0.3 mol) of 1,2,3,4-tetrahydroquinoline in 100 ml of ether. Stirring was then continued for another hour. The separated 1,2,3,4-tetrahydroquinoline hydrobromide was sucked off. The filtrate was evaporated to dryness and the resulting residue (50 g of a viscous oil) was cyclised without purification.

55

150 ml of ethanol and 81 g of anhydrous zinc chloride were added to 50 g of the crude product. This mixture was boiled, while stirring, for 16 hours. A large part of the ethanol was then distilled off. After cooling to room temperature, 200 ml of 2N HCl and 500 ml of ether were added to the residue. This mixture was stirred until nearly everything had dissolved. The aqueous layer was extracted. The collected ether layers were dried and then evaporated to dryness. The residue (17 g) was purified by chromatography over 900 g of silicagel, elution being carried out with dichloromethane. Yield 8.8 g (26% with respect to 1,2,3,4-

tetrahydroquinoline) having a melting-point of 70-72 °C.

b) 4,5,6,7-tetrahydro-pyrrolo-[3,2,1-jk]-1-benzazepin-1-carboxylic acid

21.2 g (95 mmol) of 1-(α -chloroacetyl)-2,3,4,5-tetrahydro-1-benzazepine were heated (130 °C), 19.0 g (142 mmol) of aluminium chloride were added, and the mixture was kept at this temperature until gas development almost had ceased. Water and methylene chloride were then added and the mixture was shaken. The methylene chloride layer was separated, evaporated in vacuo, and chromatographed over silicagel with ether/petroleum ether (1/1) as eluent. Yield 8.85 g (50%) of 1,2,4,5,6,7-hexahydro-pyrrolo-[3,2,1-jk]-1-benzazepin-2-one.

7.5 g (0.04 mol) of this product were dissolved in 80 ml of diglyme, 5.7 g (0.044 mol) of borotrifluoro etherate were added, and the mixture was heated at 80 °C. 1.7 g (0.044 mol) of sodium borohydride were added in small amounts, and the mixture was stirred at 80 °C for another 2 hours.

100 ml of 2N hydrochloric acid were added dropwise at 100 °C, and the mixture was poured on ice, made alkaline, extracted with ether, washed with 6N hydrochloric acid, water and a sodium bicarbonate solution, and evaporated. Yield 3.3 g (48%) of 4,5,6,7-tetrahydro-pyrrolo-[3,2,1-jk]-1-benzazepine.

2.56 g (16.7 mmol) of phosphorus oxychloride were slowly added to 20 ml of dimethylformamide under nitrogen at 10-20 °C. This mixture was cooled to 10 °C, and a solution of 2.6 g (15.2 mmol) of the obtained benzazepine in 20 ml of dimethylformamide was added dropwise. The mixture was stirred for 30 minutes at 10 °C and for 1 hour at 40 °C. The mixture was poured on ice, made alkaline with 4 ml of 50% NaOH, boiled during 10 minutes and cooled. Then the mixture was extracted with ether, dried and evaporated. Yield 2.9 g (96%) of 1-formyl-4,5,6,7-tetrahydro-pyrrolo-[3,2,1-jk]-1-benzazepine.

1.0 g (5 mmol) of this product was dissolved in 75 ml of acetone, and a solution of 0.53 g (3.7 mmol) of potassium permanganate in 25 ml of water was added. The mixture was stirred for 1 hour, and filtered on hyflo. The filtrate was evaporated in vacuo, and 300 ml of ethyl acetate were added to the residue. The mixture was acidified with concentrated hydrochloric acid while stirring firmly, dried, filtered and evaporated. Yield 1.0 g (92%), melting-point 200-201 °C.

The following compounds have been prepared in a similar way:

- (1) 5,6-dihydro-4H-pyrrolo-[3,2,1-ij]-quinolin-1-carboxylic acid; melting-point 221-223 °C;
- (2) 7-fluoro-5,6-dihydro-4H-pyrrolo-[3,2,1-ij]-quinolin-1-carboxylic acid; melting-point 231.5-232 °C;
- (3) 6,7-dihydro-indolo-[1,7-ab]-[1]-benzazepin-2-carboxylic acid.


c) 6,7-dihydro-indolo-[1,7-ab]-[1]-benzazepin-1-carboxylic acid

35 mmol (12.0 g, 60%) N-amino-iminodibenzyl and 4.5 g (38 mmol) of ethyl pyruvate were dissolved in 200 ml of alcohol and boiled for 30 minutes. Thereafter the reaction product of 15 ml of acetyl chloride and 50 ml of alcohol was added, and the mixture was boiled for 1 hour, evaporated and shaken with 2N sodium hydroxide and ethyl acetate. The ethyl acetate solution was evaporated and chromatographed over silicagel with ether/petroleum ether (1/9) as eluent. Yield 7.4 g (72%) oil.

The obtained product was boiled for 1 hour with an aqueous solution of KOH in alcohol, evaporated in vacuo; the residue was shaken with 2N hydrochloric acid and ethyl acetate. The ethyl acetate solution was washed with water, dried and evaporated. Yield 4.8 g (72%), melting-point 221-222 °C.

Claims

1. Indole derivatives of the general formula 2.



(2)

wherein

- 15 R₁ is straight or branched alkyl having 1-4 C-atoms; fluorinated or hydroxylated alkyl having 1-4 C-atoms, or R₁ is straight or branched alkenyl or alkynyl having 2-4 C-atoms, or R₁ is straight or branched alkoxy or alkylthio having 1-6 C-atoms which may be substituted with one or more fluorine atoms or hydroxyl groups, or with one optionally substituted phenyl group, or two alkoxy groups R₁ bound to adjacent carbon atoms may form a ring consisting of 5-7 ring atoms, or R₁ is straight or branched (C₁₋₄)alkoxy-, (C₁₋₄)alkylthio- or
 20 alkyl-(C₁₋₄)sulphonyl-(C₁₋₂)alkyl which may comprise one or more fluorine atoms or hydroxyl groups, or R₁ is an alkoxycarbonyl methyl group having 1-6 C-atoms in the alkoxy group, or R₁ is a group R₄R₅N-CO-CH₂-, R₄R₅N-CO- or R₄R₅N-SO₂- wherein R₄ and R₅ independently of each other are hydrogen, alkyl having 1-3 C-atoms, or together with the nitrogen atom form a heterocyclic 5- or 6-ring, or R₁ is hydroxy, halogen, cyano straight or branched alkoxycarbonyl having 1-6 C-atoms in the alkoxy group;

25 n has the value 0-3;


Z together with the carbon atom and the nitrogen atom and the intermediate carbon atom forms a heterocyclic group consisting of 5-8 ring atoms, in which, besides the nitrogen atom already present, a second hetero atom from the group N, O, S, S-O, SO₂ or C=O may be present, which ring may be substituted with 1-3 alkyl groups having 1-4 C-atoms, a phenyl group or a spiroalkyl group (C₂₋₅), or which
 30 ring may be annelated with a saturated or non-saturated carbocyclic or heterocyclic ring consisting of 5- or 6-ring atoms which may be substituted with halogen or alkyl having 1-4 C-atoms;

R₃ is hydrogen, halogen, alkyl, alkoxy or alkylthio having 1-4 C-atoms, and


R₂ is the group -B-C-D, wherein B is the group -CO-, C is oxygen or NR₆, wherein R₆ is hydrogen, (C₁₋₇) alkyl or benzyl, and D is a group (CH₂)_m-R₇, wherein m has the value 0, 1 or 2 and R₇ is a carbocyclic ring system in which one carbon atom has been replaced by a secondary or tertiary nitrogen atom or the N-oxide thereof;

and the pharmacologically acceptable acid addition salts thereof.


2. Indole derivatives as claimed in Claim 1, characterized in that D is a group of the formula 3, 4 or 5



(3)



(4)




(5)

50 wherein p has the value 1 or 2, s has the value 0, 1 or 2, q is 2, 3 or 4, r is 1, 2 or 3, and wherein R₈ is hydrogen, (C₁₋₇)alkyl, (C₃₋₈)cycloalkyl, (C₃₋₈)cycloalkyl-(C₁₋₂)alkyl, (C₃₋₅)alkenyl, (C₃₋₅)alkynyl, or wherein R₈ is a group (CH₂)_t-R₉, wherein t is 1 or 2 and R₉ is hydroxy, or R₉ is the phenyl group, thienyl group, pyrrolyl group or furyl group, which groups may be substituted with 1 or 2 substituents, and R₁₀ is absent or is an oxygen atom.

55 3. Pharmaceutical compositions, characterized in that they comprise at least one compound as claimed in Claim 1 as an active substance.

4. A method of treating symptoms which are caused by over-excitation of 5-HT receptors, characterized in that a composition as claimed in Claim 3 is used.

5. A method of preparing compounds as claimed in Claim 1, characterized in that
 a) a compound of formula 6



(6)

wherein R_1 , n , R_3 , Z and B have the meanings mentioned in Claim 1 and X is a group which may be replaced by a nucleophile, is converted with a compound of the formula $Y-C-D$ (7), wherein C and D have the meanings mentioned in Claim 1 and Y is hydrogen or an alkali metal atom; or

- b) the amino-hydrogen atom of group D in a compound of formula 2, wherein R_1 , n , R_3 , Z , B , C and D have the meanings mentioned in Claim 1, with the proviso that the ring nitrogen atom in group D is a secondary nitrogen atom, is substituted by a group R_8 , wherein R_8 has the meaning mentioned in Claim 2; or

- c) a compound of formula 2, wherein R_1 , R_2 , R_3 , n and Z have the meanings given in claim 1, on the understanding that R_{10} is an oxygen atom, is prepared by oxydation of a corresponding compound having formula 2 wherein R_{10} is lacking.



EP 88 20 2800

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
A	JOURNAL OF HETEROCYCLIC CHEMISTRY, vol. XI, June 1974, pages 387-393, Orem, Utah, US; E.A. STECK et al.: "Some 5,6-dihydro-4H-pyrrolo[3,2,1-i,j]quinolines" * Pages 387-388, table 1 * ---	1	C 07 D 519/00 C 07 D 471/06 C 07 D 487/06 A 61 K 31/46
D,A	EP-A-0 189 002 (SANDOZ) * Pages 1-5; page 19, claims * -----	1-5	
			TECHNICAL FIELDS SEARCHED (Int. Cl.4)
			C 07 D 519/00 C 07 D 471/06 C 07 D 487/06
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 23-03-1989	Examiner FRANCOIS J.C.L.
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons ----- & : member of the same patent family, corresponding document			